

# **Analysis of Fine Structures of Flows, Hydrography, and Fronts in Taiwan Strait**

PI: Quanan Zheng

Department of Atmospheric and Oceanic Science, University of Maryland, College Park, MD 20742

Tel.: (301) 405-8253 Fax: (301) 314-9482 e-mail: [quanan@atmos.umd.edu](mailto:quanan@atmos.umd.edu)

Grant Number: N00014-05-1-0606

## **LONG-TERM GOALS**

The long-term goal of the project is to determine the fine structures of flows, hydrography, and fronts by analyzing a collection of in situ data and satellite data. The in situ data provide a dual purpose: the description of conditions as they exist, and the verification and validation of the results from the satellite data analysis and dynamical analysis.

## **OBJECTIVES**

(1) To resolve the fine structures of the flow field, (2) To determine the hydrodynamics and hydrography across the fronts, and (3) To examine the effects of the Taiwan Island to the fine structures and fronts under different forcing conditions. The tasks serving these objectives include the analysis of observational data from moorings and cruises, the use of an inverse analysis and extraction of flow field using Synthetic Aperture Radar (SAR) for fine structures, and the analysis of AVHRR/MODIS SST and MODIS/SeaWiFS ocean color parameters in the identification of the fine structures.

## **APPROACH**

**1) Field data analysis.** The field data are used to describe the fine structures of flows, hydrography, and fronts, and examine the effects of the bathymetry and morphology on the fine structures to achieve the objectives. The horizontal and vertical structures, the effect of various waters of contrasting characteristics under seasonal wind conditions, the effect of Taiwan Island, and the effect of the generally shallow strait with sharp depth changes at near the northeast and southeast openings of the strait are major targets. Recently measured CTD data and historical XBT/AXBT data are used to determine the vertical thermal structure in the study area. The historical data are obtained from NOAA archives, which are in the public domain.

**2) Satellite images.** Satellite SAR is the most powerful sensor for ocean remote sensing because of its all weather, all day abilities, and high spatial resolution. The spatial resolution of the state-of-the-art satellite SAR images reaches 20 – 30 m, and the swath width reaches 100 – 450 km. These specifications match the requirements for observing the ocean internal waves with a spatial characteristic scale within a range from 0.1 km to 10 km. The SAR images used for the project were taken by the ERS-1 and 2, the RADARSAT-1 and 2 satellites.

**3) Data processing methods.** TECPLOT and GIS software packages are used for the visualization of the results from the analyses of the in situ data. These results will serve not only for the description and quantification of the fine structures, but also for the validation and verification of the analysis of

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE <b>30 SEP 2006</b>		2. REPORT TYPE		3. DATES COVERED <b>00-00-2006 to 00-00-2006</b>	
4. TITLE AND SUBTITLE <b>Analysis of Fine Structures of Flows, Hydrography, and Fronts in Taiwan Strait</b>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>University of Maryland, College Park, Department of Atmospheric and Oceanic Science, College Park, MD, 20742</b>				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release; distribution unlimited</b>					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT <b>Same as Report (SAR)</b>	18. NUMBER OF PAGES <b>6</b>	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			

satellite data including the algorithms of the inverse analysis of flow field using the satellite SAR images. Commercialized image processing software packets will be used for imagery enhancement, orthorectification, filtering, and data extraction.

**4) Theoretical analysis.** The objectives of theoretical analysis are to understand the nature, physics, mechanism, and laws of variation of the studied process, to derive unknown geophysical parameters using parameters and information derived from SAR image interpretation as inputs, to analyze the relation between the studied process and the surrounding environment, and to predict the future development of the studied process.

Dr. Quanan Zheng serves as a PI for research in University of Maryland. Zheng focuses on data collection, satellite image interpretation, and the theoretical analysis.

## **WORK COMPLETED**

**1) Data collection.** CTD data measured by 189 casts at 55 stations along 5 transects in the southwestern Taiwan Strait in summer 2005 are collected. Preliminary data analysis has been completed.


**2) Dynamical analysis.** Dynamical analysis of bottom-topography-induced stationary IW in NSCS has been completed. Nonlinear analysis of effects of shoaling thermocline on the IW generation has been completed. The role of Kuroshio playing in IW generation in the Luzon Strait has been analyzed using linear wave models and the Fourier transform methods.

**3) Paper preparation.** Based on the results, five papers (see **PUBLICATIONS**) have been prepared and submitted to international-circulated and peer-reviewed journals. Two of them are accepted and in press.

## **RESULTS**

### **1) Cruise data collected.**

The cruise was carried out by R/V Yanping 2 from July 4 to 15, 2005 [Chen et al., 2005]. As shown in Figure 1, the survey area covers a rectangular region of 130 by 160 km enclosed by lines linking four points from A1 (23.6390°N, 117.8522°E) clockwise in turn to A9 (22.5331°N, 118.9277°E), C9 (21.6730°N, 117.9168°E), and C1 (22.7979°N, 116.7755°E). The water depth varies from 25 m to 500 m. The purpose of this cruise observation is to collect the temperature and salinity profile data for understanding the three-dimensional (3-D) fine structure of the circulation. During the cruise observation, the survey area was scanned by 189 CTD casts at 55 stations along 5 transects. The horizontal resolution of observation grids is about 20 by 35 km.




**Figure 1.** A map of study area. Codes represent CTD cast stations. Numerals on isobaths are in m.

**2) Finding a low salinity tongue.** Using the cruise data and interpolation methods, 2-D distribution maps of salinity at depths of 5, 10, 15, 20, and 30 m are generated as shown in Figures 2a-e. One can see that the most remarkable features are a low salinity tongue sandwiched by high salinity waters on both shallow and deep water sides. The low salinity tongue runs southwest-northeastward along 40-50 m isobaths. The tongue axis is located at an offshore distance of about 60 km and generally parallel to the direction of coastline. Defining 32 psu isohaline as the boundary of the low salinity tongue [Wu, 1989], the maximum width of the tongue at the surface layer (depth of 5 m) reaches 70 km. The water tongue tip reaches 23.2°N, 118.3°E. The water tongue root is located on the southwest side of the survey area, where the lowest salinity is only 27 psu, implying that the low salinity water originates from somewhere southwest of the survey area.

Vertically, at the 10 m layer, the water tongue keeps patterns similar to that at the surface layer. The location of water tongue tip keeps unchanged, but the width narrows to a maximum of about 45 km at the water tongue root. The lowest salinity increases 2.5 psu to 29.5 psu. The low salinity water exists even at the bottom layer (30 m), but cannot keep an entire tongue pattern below the 15 m layer. In other words, the low salinity water is mainly concentrated in the upper layer from the sea surface to about 10 m.

Horizontally, the salinity gradients can be calculated. Taking Figure 3b (10 m layer) as an example, along the west boundary of survey area, the salinity gradients reach  $0.24 \text{ psu km}^{-1}$  on the shallow water side, and  $0.085 \text{ psu km}^{-1}$  on the deep water side, respectively. This implies the existence of strong salinity fronts on the both sides of the low salinity tongue according to a generally accepted criterion of  $0.018 \text{ psu km}^{-1}$  for a salinity front [Zhuang et al., 2003]. Meanwhile, the existence of strong salinity fronts implies that a jet-like current has to serve as low salinity water supplier to maintain the existence of low salinity tongue [Zheng et al., 2004].



***Figure 3. Two-dimensional distribution maps of salinity at depths of 5 (a), 10 (b), 15 (c), 20 (d), and 30 m (e) derived from the 2005 summer cruise data by interpolation methods.***

**3) Dynamical analysis of bottom-topography-induced stationary IW in Taiwan Strait.** The satellite SAR images display wave-like patterns of the ocean bottom topographic features at the south outlet of Taiwan Strait (TS). Field measurements indicate that the most TS water body is vertically stratified. In order to explore the mechanisms and to determine the relations between the SAR imagery and the bottom features, a two-dimensional, three-layer ocean model with sinusoidal bottom topographic features is developed. Analytical solutions and inferences of the momentum equations of the ocean model lead to the following solutions. 1). In the lower layer, the topography-induced waves (topographic waves) exist in the form of stationary waves, which satisfy a lower boundary resonance condition  $\sigma = kC_0$ , here  $\sigma$  is an angular frequency of the stationary waves,  $k$  is a wavenumber of bottom topographic corrugation, and  $C_0$  is a background current speed. 2). As internal waves, the topographic waves may propagate vertically to the upper layer with an unchanged wavenumber  $k$ , if a frequency relation  $N_3 < \sigma < N_2$  is satisfied, here  $N_2$  and  $N_3$  are the Brunt-Wäisälä frequencies of middle layer and upper layer, respectively. 3). The topographic waves are extremely amplified if an upper layer resonance condition is satisfied. The theoretical SAR image of topographic waves is derived on the basis of current-modulated small wave spectra. The results indicate that the topographic waves on SAR images have the same wavelength of bottom topographic corrugation, and the imagery brightness peaks are either inphase or antiphase with respect to the topographic corrugation, depending on a sign of a coupling factor. These theoretical predictions are verified by field observations. The results provide a physical basis for quantitative interpretation of SAR images of bottom topographic waves in the stratified ocean.

## IMPACT/APPLICATIONS

The results of this project will provide the users fine structures of ocean circulation in Taiwan Strait, and will benefit the broader oceanographic community, ocean engineering industries, underwater navigation and operational users. The results may also serve as a basis for empirical, theoretical, and numerical prediction models of ocean circulation in the Taiwan Strait. The results will further reveal SAR imaging mechanisms and be used for SAR image interpretation.

## RELATED PROJECTS

This project and the project of Grant Number N00014-05-1-0605 at the Louisiana State University (PI: Dr. Chunyan Li) are two separately funded components of the same project. The PI of this project serves as a PI for an ongoing ONR PO project titled “Satellite Synthetic Aperture Radar Detection of Ocean Internal Waves in the South China Sea”. The study areas of two projects are immediately adjacent. Therefore, two projects sometimes share the same data resources of field observations.

## REFERENCES

Chen, Z., J. Hu, B. Huang, and H. Hong, *Temperature and Salinity Data Report: Cruise in the Southwestern Taiwan Strait during July 2005*, Technical Report: No. CMI-001, Xiamen Univ., Xiamen, pp. 1-200, 2005.

Wu, B., A study on the circulation in shelf waters west to Zhujiang Estuary I. Coastal current, *J. Oceanogr. Taiwan Str.*, 8, 360-365, 1989.

Zheng, Q. and V. Klemas, Determination of winter temperature patterns, fronts, surface currents in the Yellow Sea and East China Sea from satellite imagery, *Rem. Sens. Environ.*, 12, 201-218, 1982.

Zhuang, W., J. Hu, Z. He, G. Zeng, and Z. Chen, An analysis on surface temperature and salinity from southern Taiwan Strait to Zhujiang River estuary during July-August, 2000, *J. Tropical Oceanogr.*, 22, 68-76, 2003.

## **PUBLICATIONS**

Zheng, Q., L. Li, X. Guo, Y. Ge, D. Zhu, and C. Li, SAR Imaging and hydrodynamic analysis of ocean bottom topographic waves. *J. Geophys. Res.*, 2006 [in press].

Hong, H., J. Hu, Z. Chen, Q. Zheng, and C. Li, Three-dimensional structure of a low salinity tongue in the southern Taiwan Strait observed in summer 2005. *Geophys. Res. Lett.*, 2006.

Zheng, Q., R. D. Susanto, C-R. Ho, Y. T. Song, and Q. Xu, Statistical and dynamical analyses of generation mechanisms of solitary internal waves in the northern South China Sea. *J. Geophys. Res.*, 2006.

Yuan, Y., Q. Zheng, D. Dai, F. Qiao, and J. Meng, The mechanism of internal waves in the Luzon Strait. *J. Geophys. Res.*, 2006 [in press].

Ho, C.-R., C.-Y. Lin, N.-J. Kuo, and Q. Zheng, Analyses of upper layer thickness variation in the South China Sea from satellite altimeter data and in situ measurements. *Rem. Sens. Environ.*, 2006.